

The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

Editor-in-Chief

Dan Mires

Editorial Board

Sheenan Harpaz Agricultural Research Organization
Beit Dagan, Israel

Zvi Yaron Dept. of Zoology
Tel Aviv University
Tel Aviv, Israel

Angelo Colorni National Center for Mariculture, IOLR
Eilat, Israel

Rina Chakrabarti Aqua Research Lab
Dept. of Zoology
University of Delhi

Ingrid Lupatsch Swansea University
Singleton Park, Swansea, UK

Jaap van Rijn The Hebrew University
Faculty of Agriculture
Israel

Spencer Malecha Dept. of Human Nutrition, Food
and Animal Sciences
University of Hawaii

Daniel Golani The Hebrew University of Jerusalem
Jerusalem, Israel

Emilio Tibaldi Udine University
Udine, Italy

Copy Editor

Ellen Rosenberg

Published under auspices of
**The Society of Israeli Aquaculture and
Marine Biotechnology (SIAMB),
University of Hawaii at Manoa Library**

and
**University of Hawaii Aquaculture
Program** in association with
AquacultureHub

<http://www.aquaculturehub.org>



UNIVERSITY
of HAWAII
MĀNOA
LIBRARY



AquacultureHub
educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:
Israeli Journal of Aquaculture - BAMIGDEH -
Kibbutz Ein Hamifratz, Mobile Post 25210,
ISRAEL

Phone: + 972 52 3965809

<http://siamb.org.il>

Effects of Lecithin on Cholesterol Digestibility in the Prawn, *Artemesia longinaris* (Crustacea, Penaeidae)

Nora S. Haran^{1*} and Jorge L. Fenucci^{1, 2}

¹ Departamento de Ciencias Marinas, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Funes 3350 (7600) Mar del Plata, Argentina

² Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina

(Received 30.7.07, Accepted 3.9.07)

Key words: *Artemesia longinaris*, Crustacea, cholesterol, lecithin, digestibility

Abstract

Cholesterol and dietary phospholipids are important to the growth and survival of crustaceans. Soy lecithin enhances cholesterol solubilization while phosphatidylcholine transfers cholesterol from the hepatopancreas to the hemolymph. To study the influence of dietary lecithin on digestibility and transport of cholesterol to the hemolymph in *Artemesia longinaris*, diets with different levels of supplemental cholesterol (0.5, 1, 1.5, and 2%) and soy lecithin (0, 0.5, 1, and 2%) were fed to prawns for two weeks. The diet containing the lowest level of cholesterol (0.5%) and lecithin (1%) resulted in the lowest apparent digestibility of cholesterol. The diets containing 2% cholesterol and 1% or 2% lecithin resulted in the highest hemolymph cholesterol contents. There was a linear relationship ($r = 0.832$; $p < 0.01$) between hemolymph cholesterol and the amount of lecithin in the diet. There were no significant differences between treatments in cholesterol contents in the hepatopancreas. In a second 6-week experiment, diets containing 1.5% cholesterol significantly improved weight gains, regardless of the lecithin content. Survival rates did not differ among treatments. Hence, the efficacy of cholesterol does not seem to be related to dietary inclusion of lecithin.

Introduction

Sterols are essential nutrients for prawns because crustaceans are unable to synthesize these compounds (Zandee, 1966; Teshima and Kanazawa, 1971). Feeding experiments have demonstrated the necessity of dietary sterols for growth and survival (Castell et al., 1975; Petriella et al. 1984).

Essential components such as molting hormones, sex hormones, bile acids, and vitamin D are synthesized from cholesterol. Cholesterol is also a membrane component and functions in absorption and transport of fatty acids (Akiyama et al., 1991). Cholesterol is obtained directly from the diet or via the meta-

* Corresponding author. E-mail: nsharan@mdp.edu.ar

bolic conversion of other dietary sterols. Meals and invertebrate marine oils are excellent sources of cholesterol (Akiyama, 1992).

Phospholipids are cell membrane components that maintain their fluidity and flexibility, and play a role in lipid digestion and absorption (Akiyama et al., 1991). Dietary phospholipids apparently assist in the mobilization of cholesterol and triglycerides from the hepatopancreas to the hemolymph (D'Abramo et al., 1982). Dietary phospholipids have positive effects on weight gain and survival, in both juveniles and shrimp larvae (Piedad-Pascual, 1986; Chen and Jenn, 1991; Gonzalez-Felix et al., 2002).

The commercially valuable prawn, *Artemesia longinaris*, is distributed along the coastal waters of Argentina, Uruguay, and Brazil from 22°30'S to 43°20'S (Boschi and Scelzo, 1968). The greatest abundance of this species occurs in Mar del Plata and Rawson, Argentina. The aims of this work were to determine the effects of soy lecithin, a phospholipid, on cholesterol digestibility and the transport of cholesterol to the hemolymph in *A. longinaris* and to study the effects of different combinations of dietary cholesterol and lecithin on growth and survival.

Materials and Methods

In the first experiment, a 2-week trial was conducted to study the effect of soy lecithin on cholesterol digestibility. Six isoproteic feeds were prepared from basal diet E with a proximal composition of 56.2% protein, 6.9% lipids, and 15.55% ash (Fenucci et al., 1983). Chromic oxide was added as an inert marker to measure apparent digestibility. The feeds contained 0.5, 1.0, 1.5, or 2.0% cholesterol and 0, 0.5, 1.0, or 2.0% soy lecithin and were prepared by the cold extrusion method (Fenucci, 1981; Table 1).

Prawns (1.35-1.45 g) were randomly distributed into 150-l glass aquaria with under-gravel filters at a density of 5 prawns/aquarium and the six diets were tested in triplicate. Prawns were acclimated for seven days during which they were fed the experimental feeds *ad libitum* daily. Water ranged 19-22°C and salinity was 33 ppt. After seven days of

acclimation, prawns were fed the experimental diets for another two weeks.

Feces were collected daily with a hand net, washed with distilled water, frozen, and dried at 45°C. Hemolymph samples were taken by syringe from all prawns in the intermolt stage, from the base of the fourth pereopod. The hepatopancreas was dissected and frozen at -20°C for subsequent laboratory analysis. Molt cycle stages were determined by observation of the uropod setae development (Petriella, 1984). Cholesterol (Aikel, 1977) and chromium oxide (Fenucci, 1981) in the feces and diets were analyzed. Cholesterol was quantified as mg/ml in the hemolymph and as a percentage of the dry weight of the hepatopancreas. Apparent cholesterol digestibility (Ap Dig) was calculated according to the formula (Fenucci, 1981): $Ap\ Dig\ (\%) = 1 - (\%Cr_2O_3\ in\ diet / 100\% Cr_2O_3\ in\ feces) \times (\% cholesterol\ in\ feces / \% cholesterol\ in\ diet) \times 100$.

A second 45-day experiment was carried out to study the combined effect of cholesterol and soy lecithin on growth and survival. Four isoproteic test feeds were prepared from the standard rations used in the first experiment, with all combinations of soy lecithin and cholesterol at levels of 0 and 1.5%.

Seven prawns (1.39-1.50 g initial wt) were stocked in under-gravel filter aquaria. Each diet was tested in triplicate. The prawns were fed daily, initially *ad libitum* and then according to their requirements. Uneaten food, dead prawns, and exuviae were removed daily before feeding. Individuals were weighed to the nearest 0.01 g prior to stocking and at the end of the experiment. To prevent mortality, they were not weighed during the trial.

Data were analyzed using Bartlett's test to determine homoscedasticity of variances, factorial ANOVA, and chi square (Sokal and Rohlf, 1979).

Results

Cholesterol contents in the diets, hemolymph, and hepatopancreas are given in Table 2. Cholesterol levels in the hemolymph were significantly higher in prawns fed C₂L₂ or C₂L₁ ($p < 0.01$). Cholesterol levels in the hepatopan-

Table 1. Composition (%) of diets with different levels of lecithin and cholesterol, fed to prawns (*Artemesia longinaris*) of 1.35-1.45 g for two weeks.

	Diet name					
	<i>C</i> _{0.5} <i>L</i> ₁	<i>C</i> _{1.5} <i>L</i> ₀	<i>C</i> ₂ <i>L</i> _{0.5}	<i>C</i> ₂ <i>L</i> ₁	<i>C</i> ₂ <i>L</i> ₂	<i>C</i> ₁ <i>L</i> ₁
Cholesterol	0.5	1.5	2	2	2	1
Lecithin	1	0	0.5	1	2	1
Wheat bran	20.95	20.95	19.95	19.45	18.45	20.45
Mussel meal			30			
Squid meal			15			
Soybean meal			5			
Fish meal			20			
Sodium hexametaphosphate			1			
Alginate			2			
Fish soluble			2			
Fish oil			1			
Chromium oxide			1			
Vitamin mix			0.55			

Table 2. Cholesterol in diets, hemolymph, and hepatopancreas. Cholesterol digestibility (%) and digested cholesterol in 100 g of diet.

Diet	In diets (%)	In hemolymph (mg/ml)	In hepatopancreas (%)	Cholesterol digestibility (%)	Digested cholesterol (g per 100 g diet)
<i>C</i> _{0.5} <i>L</i> ₁	0.83	0.166±0.0290	0.774±0.0757	59.55±2.192**	0.49±0.018
<i>C</i> ₁ <i>L</i> ₁	1.48	0.122±0.0262	0.554±0.0413	85.32±0.990	0.71±0.008
<i>C</i> _{1.5} <i>L</i> ₀	1.96	0.139±0.0133	0.799±0.0870	84.28±0.445	1.65±0.008
<i>C</i> ₂ <i>L</i> _{0.5}	2.46	0.115±0.0171	1.083±0.1287	84.90±0.714	2.08±0.021
<i>C</i> ₂ <i>L</i> ₁	2.50	0.207±0.0265*	0.740±0.0865	80.48±1.520	2.01±0.038
<i>C</i> ₂ <i>L</i> ₂	2.38	0.235±0.0122*	1.362±0.1983	79.17±0.898	1.88±0.028

* significantly different at $p < 0.01$ ** significantly different at $p < 0.05$

creas did not differ significantly ($p > 0.05$) between diets. The diet with the lowest cholesterol digestibility was diet *C*_{0.5}*L*₁, which contained the smallest percentage of cholesterol. There were no significant differences in appar-

ent digestibility among the other diets ($p > 0.05$). Cholesterol digestibility was adjusted to a second order polynomial curve (Fig. 1). There was a significant correlation ($r = 0.832$; $p < 0.01$) between cholesterol in the hemolymph and




Fig. 1. Relationship between apparent digestibility of cholesterol in prawns (*Artemesia longinaris*) and the percent of cholesterol in the diet.




Fig. 2. Relationship between cholesterol in the hemolymph of prawns (*Artemesia longinaris*) and the percent of supplemental lecithin in the diet.

total lecithin in the diet (Fig. 2).

Cholesterol supplementation significantly improved growth ($p < 0.01$) while lecithin supplementation did not (Table 3). Survival was not affected by the addition of cholesterol or soy lecithin ($p > 0.05$). There was no significant ($p > 0.05$) interaction between dietary lecithin and cholesterol on growth or survival.

Discussion

The present work shows that, in *A. longinaris*, the cholesterol concentration in the hemolymph is related to dietary lecithin. Phospholipids play an important role in the metabolism of crustacea. In the hemolymph of *Marsupenaeus japonicus*, the transport of dietary lipids is carried out mainly by phospholipids associated with high density lipoproteins (Kanazawa, 1981). Crustacean hemolymph has a lower enzymatic activity than mammalian blood (Mankura et al., 1980). In mammals, dietary cholesterol enters the circulatory system after being esterified by the action of lecithin-cholesterolacyltransferase which catalyzes the transference of an acyl group from the β -position of phosphatidylcholine to the β -hydroxy group of cholesterol (Teshima, 1997).

Phosphatidylcholine is the active compo-

nent of soy lecithin in purified diets (D'Abramo et al., 1981). In *Homarus* sp., the lack of a dietary source of phosphatidylcholine apparently results in a phosphatidylcholine deficiency in the hemolymph that limits the effective cholesterol transport from the hepatopancreas to the hemolymph (D'Abramo et al., 1982). The beneficial effect of dietary phospholipids in *M. japonicus* might be due to better mobilization of lipids from the hepatopancreas or gut to the hemolymph, enhancing lipid deposition in tissues and increasing energy available for growth (Kontara et al., 1998). The absence of dietary soy lecithin can cause failure of juvenile lobsters to extricate themselves from old exoskeletons during molt (Conklin et al., 1980) while a lack of phosphatidylcholine can cause death during molt in *Homarus* sp. (D'Abramo et al., 1982).

There was no interaction between dietary lecithin and cholesterol on growth and survival. In the shrimp, *Fenneropenaeus penicillatus*, diets supplemented with phosphatidylcholine or cholesterol significantly improved weight gain but the interaction between the phosphatidylcholine and the cholesterol did not cause a significant effect on growth, food conversion, or survival (Chen and Jenn, 1991). Likewise, there were no significant

Table 3. Mean weight and survival in prawns (*Artemesia longinaris*; 1.39-1.50 g) fed diets containing different amounts of cholesterol and lecithin for 45 days.

Diet	Mean wt			Survival (%)
	Initial (g)	Final (g)	Change (%)	
C ₀ L ₀	1.39±0.348	1.73±0.318 ^a	24.46	90.48
C ₀ L _{1.5}	1.40±0.288	1.75±0.273 ^a	25.00	95.24
C _{1.5} L ₀	1.39±0.263	1.96±0.366 ^b	41.01	100.00
C _{1.5} L _{1.5}	1.42±0.233	1.98±0.276 ^b	39.44	90.48

Different superscripts in a column indicate significant differences ($p < 0.01$).

effects from the interaction of lecithin and cholesterol on growth and survival of *Penaeus monodon* zoea, mysis, and postlarvae (Paibulkichakul et al., 1998). On the other hand, there was an interaction between dietary cholesterol and phospholipids that affects growth in *L. vannamei* postlarvae (Emery, 1987, in Castille et al., 2004).

There was no significant effect of dietary lecithin on survival in *A. longinaris*. Likewise, there were no significant effects of lecithin on growth and survival of *H. americanus* fed different combinations of lecithin and cholesterol (Kean et al., 1985) or on final weight, percentage weight gain, or growth rate of juvenile *Cherax quadricarinatus* fed diets containing increasing percentages of soy lecithin for 10 weeks (Thompson et al., 2003). The highest weight gain, survival, and feed efficiency in *Fenneropenaeus merguensis* were obtained in the group fed diets containing 1-2% lecithin but there was no advantage to adding cholesterol (Thongrod and Boonyaratpalin, 1998).

On the other hand, the best apparent digestibility of cholesterol was obtained with the diet containing 2.46% cholesterol. The optimum percentage of dietary cholesterol varies. Kanazawa et al. (1971) reported that 0.5% dietary cholesterol promotes good growth in *M. japonicus*, whereas Deshimaru and Kuroki (1974) obtained the best growth with 2.1% dietary cholesterol. Kanazawa (2001) determined that the dietary cholesterol

requirements in crustaceans ranges 0.1-2.0% of the dry weight of the diet. Better survival was obtained in *A. longinaris* fed a defatted diet with added cholesterol than a diet lacking cholesterol (Petriella et al., 1984). In the same species, best growth and survival was obtained with a diet containing 2.5% cholesterol while optimum digestibility was obtained with 2.3% cholesterol (Martinez Romero et al., 1991).

Nutrient requirements vary with the size of the prawn. The assimilation pattern of proteins in small juvenile *Litopenaeus stylirostris* (1-4 g) differs from that in large animals while a positive linear relation was determined between carbohydrate assimilation and mean weight in juvenile *L. setiferus* (Fenucci, 1981). The cholesterol requirement in juvenile *L. vannamei* was estimated to be 0.35% in the absence of supplemental phospholipids but only 0.14% and 0.13%, in diets containing 1.5% and 3% phospholipids, respectively. When 5% phospholipids were provided, 0.05% dietary cholesterol was needed for optimal growth (Gong et al., 2000). Indeed, recommended cholesterol levels in commercial shrimp feeds vary 0.40-0.25% in animals of 0.5-40.0 g, i.e., a higher level of cholesterol is necessary for smaller *L. vannamei* (Akiyama et al., 1991). A change in the digestion ability with the size of the shrimp was suggested as the explanation.

While the function of phospholipids in cho-

lesterol transport is well known, further studies are needed to determine possible effects of interactions between the two compounds on the growth and survival of different sized prawns.

Acknowledgements

This work was founded by Grants PIP 3109 and 2902/03 from CONICET, Argentina.

References

- Aikel F.**, 1977. *Manual de Analisis Clinicos*. Editorial Medica Panamericana, Buenos Aires. 529 pp.
- Akiyama D.**, 1992. Future considerations for shrimp nutrition and aquaculture feed industry. pp. 198-204. In: J. Wyban (ed.). *Proc. Special Session on Shrimp Farming*. World Aquac. Soc., Baton Rouge. 301 pp.
- Akiyama D., Dominy W. and A. Lawrence**, 1991. Penaeid shrimp nutrition for the commercial feed industry: Revised. pp. 80-89. In: Akiyama, Tan (eds.). *Proc. Aquaculture Feed Processing and Nutrition Workshop*. Thailand and Indonesia. 241 pp.
- Boschi E.E. and M. Scelzo**, 1968. Nuevas campañas exploratorias camarónicas en el litoral argentino 1967/1968. *CARPAS* 4, 9:1-6.
- Castell J. E. Mason and J. Covey**, 1975. Cholesterol requirements of juvenile American lobster (*Homarus americanus*). *J. Fish. Res. Can.*, 32(8):1431-1435.
- Castille F., Lawrence A., Buisman P. and R. Drost**, 2004. Effects of sterol supplements (cholesterol FG, cholesterol SF, and sterols M1M) on growth and survival of the shrimp, *Litopenaeus vannamei* Boone. pp. 504-517. In: L. Cruz-Suarez, D. Ricque Marie, M. Nieto Lopez, D. Villarreal, U. Scholz, M. Gonzalez (eds.). *Avances en Nutricion Acuicola VII. Memorias del VII Simposio Internacional de Nutricion Acuicola*. November 16-19, 2004. Hermosillo, Sonora. 592 pp.
- Chen H and J. Jenn**, 1991. Combined effects of dietary phosphatidylcholine and cholesterol on the growth, survival and body lipid composition of marine shrimp, *Penaeus penicillatus*. *Aquaculture*, 96:167-178.
- Conklin D., D'Abramo L., Bordner C. and N. Baum**, 1980. A successful diet for the culture of juvenile lobster: The effect of lecithin. *Aquaculture*, 21:243-249.
- D'Abramo L., Bordner C., Conklin D. and N. Baum**, 1981. Essentiality of dietary phosphatidylcholine for the survival of juvenile lobsters. *J. Nutr.*, 111:63-69.
- D'Abramo L., Bordner C. and D. Conklin**, 1982. Relationship between dietary phosphatidylcholine and serum cholesterol in the lobster *Homarus sp.* *Mar. Biol.*, 67:231-235.
- Deshimaru O. and K. Kuroki**, 1974. Studies on a purified diet for a prawn. II. Optimum content of cholesterol and glucosamine in the diet. *Bull. Jap. Soc. Sci. Fish.*, 49(4):421-424.
- Fenucci J.**, 1981. *Studies on the Nutrition on Marine Shrimp of Genus Penaeus*. Ph.D. Diss., Faculty Dept. Biology, Univ. Houston, USA. 185 pp.
- Fenucci, J., Petriella A. and M. Muller**, 1983. Estudios sobre el crecimiento del camarón *Artemesia longinaris* alimentados con dietas preparadas. *Contrib. INIDEP* 424, Mar del Plata. 15 pp.
- Gong H., Lawrence A.L., Jiang D.H., Castille F.L. and D.M. Gatlin III**, 2000. Lipid nutrition of juvenile *Litopenaeus vannamei*. I. Dietary cholesterol and de-oiled soy lecithin requirements and their interaction. *Aquaculture*, 190:305-324.
- Gonzalez-Felix, M.L., Lawrence A.L., Gatlin III D.M. and M. Perez-Velazquez**, 2002. Growth, survival and fatty acid composition of juvenile *Litopenaeus vannamei* fed different oils in the presence and absence of phospholipids. *Aquaculture*, 205:325-343.
- Kanazawa A.**, 1981. Penaeid Nutrition. pp. 87-105. In: Pruder, Langdon, Conklin (eds.). *Proc. 2nd. Int. Conf. Aquac. Nutr.: Biochemical and Physiological Approaches to Shellfish Nutrition*. Spec. Publ. 2, Baton Rouge. 444 pp.
- Kanazawa A.**, 2001. Sterols in marine invertebrates. *Fish. Sci.*, 67:997-1007.
- Kanazawa A., Tanaka N., Teshima S. and K. Kashiwada**, 1971. Nutritional requirements of prawn. II: Requirements for sterols. *Bull. Jap. Soc. Sci. Fish.*, 37: 211-215.
- Kean J.C., Castell J.D., Boghen A.G., D'Abramo L.R. and D.E Conklin**, 1985. An evaluation of the lecithin and cholesterol

- requirement of juvenile lobster (*Homarus americanus*) using crab protein-based diets. *Aquaculture*, 47:143-149.
- Kontara E., Djunaidah I., Coutteau P. and P. Sorgeloos**, 1998. Comparison of native, lyso and hydrogenated soybean phosphatidylcholine as source of phospholipids in the diet of postlarval *P. japonicus*. *Arch. Anim. Nutr.*, 51:1-19.
- Mankura M., Dalimunthe D. and M. Kayama**, 1980. Comparative biochemical studies on plasma cholesterol. II. Relationship between plasma esterified cholesterol and lecithin: Cholesterol acyltransferase activity. *Bull. Jap. Soc. Sci. Fish.*, 46(5):583-586.
- Martinez Romero P., Casal de Fenucci A. and J. Fenucci**, 1991. Dietary cholesterol influence on the growth and survival of the Argentine prawn *Artemesia longinaris* Bate. *J. Aquac. Trop.*, 6:111-117.
- Paibulkichakul C., Piyatiratitivorakul S., Kittakoop P., Viyakarn V., Fast A. and P. Menasveta**, 1998. Optimal dietary levels of lecithin and cholesterol for black tiger prawn *Penaeus monodon* larvae and postlarvae. *Aquaculture* 167: 273-281.
- Petriella A.M.**, 1984. Estudio del ciclo de muda del camaron *Artemesia longinaris* (Decapoda, Penaeidae) Bate. I. Setogenesis. *Physis, Secc. A*, 42(103):93-100.
- Petriella A.M., Muller M.I., Fenucci J.L. and M.B. Saez**, 1984. Influence of dietary fatty acids on the growth and survival of the Argentine prawn *Artemesia longinaris* Bate. *Aquaculture*, 37:11-20.
- Piedad-Pascual F.**, 1986. Effect of supplemental lecithin and lipid sources on the growth and survival of *Penaeus monodon* juveniles. pp. 615-618. In: J.L. Maclean, L.B. Dizon, L.V. Hosillos (eds.). *Proc. 1st Asian Fisheries Forum*. Asia Fisheries Society, Manila, Philippines.
- Sokal R. and F. Rohlf**, 1979. *Biometria*. Madrid. 835 pp.
- Teshima S.**, 1997. Phospholipids and sterols. pp. 85-107. In: L.R. D'Abramo, D.E. Conklin, D.M. Akiyama (eds.). *Advances in World Aquaculture, Vol. 6. Crustacean Nutrition*. World Aquac. Soc., Baton Rouge. 587 pp.
- Teshima S. and A. Kanazawa**, 1971. Biosynthesis of sterols in the lobster, *Panulirus japonica*, the prawn, *Penaeus japonicus*, and the crab, *Portunus trituberculatus*. *Comp. Biochem. Physiol.*, 38B:597-602.
- Thongrod S. and M. Boonyaratpalin**, 1998. Cholesterol and lecithin requirement of juvenile banana shrimp, *Penaeus merguensis*. *Aquaculture*, 161:315-321.
- Thompson K., Muzinic A., Christian T., Webster C., Manomaitis L. and D. Rouse**, 2003. Lecithin requirements of juvenile Australian red claw crayfish *Cherax quadricarinatus*. *Aquac. Nutr.*, 9(4):223-230.
- Zandee D.**, 1966. Metabolism in the crayfish *Astacus astacus* (L). III. Absence of cholesterol synthesis. *Physiol. Biochem.*, 74(3):435-441.