

Original Research Articles

The Thermal Effects of Coastal Nuclear Power Plant Thermal Discharge on Two Fish Species Based on the Probabilistic Frequency of Temperature Increment

Lei Li^{1a}, Chenshan Shao^{2b}, Jiaying Cai², Baojun Tang³, Weiwei Su², Weiyi Zou², Mei Jiang^{4c}¹ East China Sea Fisheries Research Institute, ² College of Marine and Biological Engineering, Yancheng Institute of Technology, ³ College of Marine and Biological Engineering, Yancheng Institute of Technology, ⁴ East China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Shanghai, China

Keywords: Thermal discharge, probabilistic frequency, fish, thermal effect

<https://doi.org/10.46989/001c.123069>

Israeli Journal of Aquaculture - Bamidgheh

Vol. 76, Issue 3, 2024

This study selected *Trachinotus ovatus* and *Nibea albiflora* as the focal subjects to investigate the impacts of probabilistic frequency of temperature increment through laboratory simulations that mimicked the seasonal water elevated temperature adjacent to two coastal nuclear power plants. This research was underpinned by the probabilistic analysis of temperature increase frequencies, aiming to evaluate the thermal tolerance of these species across different seasonal contexts. Results demonstrated that the *T. ovatus* did not exhibit mortality in the spring, autumn, and winter. Mortality in *T. ovatus* occurred only during the summer in temperature-duration probability experimental groups subjected to 8.5°C above ambient temperature with exposure probabilities of 50%, 75%, and 100%. The average survival rates were (93.3±3.3) %, (93.3±5.1) %, and (86.7±2.9) % respectively. The *N. albiflora* experienced mortality in spring, summer, autumn, and winter, with the highest mortality rate in summer. Compared to *N. albiflora*, *T. ovatus* exhibited significantly higher thermal tolerance across all seasons. As the experimental temperature increased and the duration probability rose, the mortality rates of both fish species showed an upward trend. There was an interaction between elevated temperature and duration probability, with the duration of different elevated temperature amplitudes significantly impacting the fish's survival rates.

INTRODUCTION

Nuclear power is one of the most important sources of electricity generation globally, accounting for approximately 14% of global electricity. It does not produce carbon emissions or other gases that affect climate change and can provide a stable supply of electricity. Therefore, nuclear power has become one of the primary sources of global electricity supply, with 448 nuclear reactors currently in operation.¹ Water is an indispensable component in the operation of nuclear power, as the steam-water cycle during nuclear power operation requires a significant amount of water. The temperature at the nuclear power plant's water discharge is about 7-10°C higher than at the intake.^{3,4} Consequently, nuclear power plants are generally located by the sea or in inland areas rich in water resources. Inevitably, the thermal discharge can have certain impacts on aquatic ecosys-

tems, including water, sediment, and marine life. Fishery resources are an essential component of natural resources, serving not only as a significant source of food for humans but also providing employment, economic benefits, and social welfare for those engaged in fishing activities.⁵ Water temperature directly affects the body temperature of most fish species. Therefore, the impact of thermal discharge on fish has been a focal point of research. Current studies on the effects of thermal discharge on fish primarily concentrate on areas such as temperature tolerance,^{6,7} growth, and development,⁸ spatial distribution,⁹ habitats and physiological ecology.⁸ Research findings indicate that the impact of thermal discharge on fish varies. Current studies generally expose fish to lethal or sublethal temperatures under thermal shock conditions. However, in the actual operation of nuclear power plants, the temperature of thermal discharge is influenced by hydrological conditions such as tides in different regions, resulting in variations in temper-

a Lei Li and Chenshan Shao contributed equally to the study

b Lei Li and Chenshan Shao contributed equally to the study

c Corresponding author Mei Jiang, Email: jiangrose73@163.com

ature over time within mixing zones. Therefore, the temperature settings of current studies are idealized, overlooking the effects of hydrological conditions like tides on the duration of exposure in different elevated temperature mixing zones, leading to discrepancies with real-world scenarios. Research in this area is lacking, and it is necessary to conduct further studies to address this gap.

The *Trachinotus ovatus* belongs to the order Perciformes, family Carangidae, and genus *Trachinotus*. It is a warm-water pelagic fish predominantly found in the tropical and temperate regions of the southern Chinese seas.⁹ The *Nibea albiflora* is part of the order Perciformes, family Sciaenidae, and genus *Nibea*. It is a demersal fish found in shallow seas, primarily distributed in China's temperate and subtropical nearshore waters.¹⁰ Both fish species are important targets for fishing and aquaculture, possessing significant economic value. Coastal waters serve as crucial areas for spawning, fattening, and foraging.

This study employed *T. ovatus* and *N. albiflora* as experimental subjects and devised a methodology predicated on the probabilistic frequency of temperature increment. Indoor experiments were conducted to emulate the operating conditions of thermal discharge from two nuclear power plants in Zhejiang and Guangdong provinces, China, across different seasons. The thermal effects of thermal discharge on the *T. ovatus* and *N. albiflora* were investigated. The findings contribute foundational data and technical insights for the construction of a more scientific impact assessment indicator system for thermal discharge.

MATERIALS AND METHODS

EXPERIMENTAL MATERIALS

Experiments on *T. ovatus* and *N. albiflora* were conducted in May 2022 at the South China Sea Fisheries Research Institute's experimental base in Shenzhen, Guangdong, and at the East China Sea Fisheries Research Institute's experimental base in Ninghai, Zhejiang, both under the Chinese Academy of Fishery Sciences. The two respective experimental bases provided the experimental fish. The *T. ovatus* had an average body length of (6.57±0.19) cm and an average body mass of (8.93±0.62) g; the *N. albiflora* had an average body length of (5.79±0.38) cm, and an average body mass of the experimental seawater was sourced from the vicinity of the two experimental bases, and was used after sedimentation, sand filtration, and disinfection, with water quality indicators being simultaneously tested. At the Shenzhen experimental base, the pH was 8.02. Salinity was 31.34, COD content was 1.23 mg·L⁻¹, ammonia nitrogen content was 0.021 mg·L⁻¹, nitrate nitrogen content was 0.005 mg·L⁻¹, nitrite nitrogen content was 0.198 mg·L⁻¹, and reactive phosphate content was 0.020 mg·L⁻¹. The test results indicated that the inorganic nitrogen (ammonia nitrogen + nitrate nitrogen + nitrite nitrogen) at the Shenzhen experimental base met the Class II standards (0.30 mg·L⁻¹) of the "Seawater Quality Standards"(GB 3097-1997); reactive phosphate also complied with the Class II standards (0.030 mg·L⁻¹) of the "Seawater Quality

Standards" (GB 3097-1997). At the Ninghai experimental base, the pH was 8.12. Salinity was 23.56, COD content was 0.54 mg·L⁻¹, ammonia nitrogen content was 0.020 mg·L⁻¹, nitrate nitrogen content was 0.457 mg·L⁻¹, nitrite nitrogen content was 0.016 mg·L⁻¹, and reactive phosphate content was 0.041 mg·L⁻¹. The test results showed that the inorganic nitrogen (ammonia nitrogen + nitrate nitrogen + nitrite nitrogen) at the Ninghai experimental base met the Class III standards (0.50 mg/L) of the "Seawater Quality Standards" (GB 3097-1997); reactive phosphate also complied with the Class III standards (0.045 mg/L) of the "Seawater Quality Standards"(GB 3097-1997).

EXPERIMENTAL METHODS

ACCLIMATION

Initially, specimens of *T. ovatus* and *N. albiflora* were cultured separately in aquaculture pools at two experimental sites. According to the surface seawater temperature conditions outside the thermal discharge mixing zones of two coastal nuclear power plants in Zhejiang and Guangdong provinces during the four seasons of spring, summer, autumn, and winter, four groups of acclimation temperatures were set. Specifically, the surface seawater temperatures near the Guangdong nuclear power plant were 21.5°C (average spring temperature), 27.5°C (average summer temperature), 25.5°C (average autumn temperature), and 17.0°C (average winter temperature); near the Zhejiang nuclear power plant, the temperatures were 14.5°C (average spring temperature), 25.0°C (average summer temperature), 22.5°C (average autumn temperature), and 10.0°C (average winter temperature). The water temperature was adjusted to each acclimation temperature at 1°C·d⁻¹ using a heating and cooling unit. The acclimation period lasted for 7 days, during which artificial feed was provided once in the morning and evening (09:00 and 17:00), with uneaten feed and feces removed, and one-third of the seawater changed after 30 minutes of feeding. An oxygen pump was continuously used throughout the acclimation process to maintain dissolved oxygen levels above 6.0 mg·L⁻¹ under natural lighting with a light cycle of 12 hours light (L) and 12 hours dark (D). After acclimation, healthy and similarly sized individuals were selected for the experiments.

EXPERIMENT ON THE IMPACT OF THERMAL SHOCK ON FISH SURVIVAL RATES

The definition of probabilistic frequency of temperature increment is the proportion of time a specific temperature increase persists within 24 hours. Based on the thermal discharge conditions and hydrological characteristics of the sea areas near two nuclear power plants in Zhejiang and Guangdong provinces, the temperature increase amplitudes were set at 0.5°C, 1.0°C, 2.0°C, 4.0°C, and 8.5°C. The proportions of the time this temperature increment persisted within 24 hours were set at 25%, 50%, 75%, and 100%, respectively.

The experiment assessing the effects of thermal shock on fish survival was conducted in 100 L glass tanks filled




Fig. 1. Survival rate of *T. ovatus* at different elevated temperature amplitudes and duration probabilities in spring

with 80 L of water. Initially, the glass tanks were submerged in larger aquaculture pools to equalize water levels, and a heating and cooling unit was used to control the water temperature. The experiment was divided into elevated and control temperature groups, with three replicates for each temperature setting. Temperature increments were regulated to within $\pm 0.5^\circ\text{C}$ using heating rods. Upon reaching the designated rise amplitude, 30 fish were introduced into each tank. The heating was stopped when the experiment reached the predetermined duration, with a total experimental time of 24 hours. For ease of description, this is called the elevated temperature-duration probability experimental group. Before the experiment, the test organisms were fasted for 24 hours. During the experiment, an oxygen pump was used to aerate the water, maintaining dissolved oxygen levels above $6.0 \text{ mg}\cdot\text{L}^{-1}$. Water temperature was monitored using a multiparameter water quality analyzer (YSI 650 MDS), and observations and records of the test fish's survival, stress, and mortality were made. The definition of fish mortality was the cessation of gill cover movement or no response to touch. Natural lighting was used throughout the experiment, with a light cycle of 12 hours light (L) and 12 hours dark (D). It was ensured that the mortality rate in the control group was $\leq 10\%$ by the end of the experiment; otherwise, the experiment would be conducted again.

DATA PROCESSING AND ANALYSIS

The statistical results are presented as mean \pm standard deviation (Mean \pm SD). Data were statistically analyzed using IBM SPSS Statistics 22.0 (IBM Inc., USA). Initially, the experimental data underwent normality (Kolmogorov-

Smirnov test) and homogeneity of variance testing (Levene test). A two-factor analysis of variance (one-way ANOVA) was used to analyze the effects of elevated temperature amplitudes and duration probabilities on survival rates and their interaction, with a significance level set at 0.05.

RESULTS

THE IMPACT OF PROBABILISTIC FREQUENCY OF TEMPERATURE INCREMENT ON THE SURVIVAL RATE OF *T. OVATUS* ACROSS DIFFERENT SEASONS

The experimental results on the impact of probabilistic frequency of temperature increment on the survival rate of *T. ovatus* across different seasons showed that under spring temperature conditions, no mortality was observed in the control group or any of the experimental groups of *T. ovatus* (Figure 1). No mortality occurred in the control group of *T. ovatus* in summer temperature conditions. However, in the experimental groups with $8.5^\circ\text{C}\text{-}50\%$, $8.5^\circ\text{C}\text{-}75\%$, and $8.5^\circ\text{C}\text{-}100\%$ elevated temperature-duration probability, mortality was observed in *T. ovatus*, with average survival rates of $(93.3\pm 3.3)\%$, $(93.3\pm 5.1)\%$, and $(86.7\pm 2.9)\%$ respectively. Among these, the $8.5^\circ\text{C}\text{-}100\%$ elevated temperature-duration probability group exhibited the highest mortality rate in *T. ovatus*, and there was a significant difference compared to the $8.5^\circ\text{C}\text{-}50\%$ and $8.5^\circ\text{C}\text{-}75\%$ elevated temperature-duration probability groups. Under autumn and winter temperature conditions, no mortality was observed in the control group or any of the experimental groups of *T. ovatus* (Figures 3-4).




Fig. 2. Survival rate of *T. ovatus* at different elevated temperature amplitudes and duration probability in summer

Note: The different letters on the same list mean significant difference ($P < 0.05$), the same below.




Fig. 3. Survival rate of *T. ovatus* at different elevated temperature amplitudes and duration probability in autumn

THE IMPACT OF PROBABILISTIC FREQUENCY OF TEMPERATURE INCREMENT ON THE SURVIVAL RATE OF *N. ALBIFLORA* ACROSS DIFFERENT SEASONS

The experimental results on the impact of probabilistic frequency of temperature increment on the survival rate of *N. albiflora* across different seasons showed that under spring temperature conditions, certain mortality was observed in

the experimental groups with 1.0°C-100%, 2.0°C-75%, 2.0°C-100%, 4.0°C-75%, 4.0°C-100%, 8.5°C-75%, and 8.5°C-100% elevated temperature-duration probability, with average survival rates of (94.4±3.8)%, (94.4±3.3)%, (94.4±2.9)%, (93.3±5.1)%, (88.9±2.3)%, (85.7±3.3)%, and (83.3±3.8)% respectively. No mortality occurred in the control group or the other elevated temperature-duration probability experimental groups of *N. albiflora*. Overall, as




Fig. 4. Survival rate of *T. ovatus* at different elevated temperature amplitudes and duration probability in winter




Fig. 5. Survival rate of *N. albiflora* at different elevated temperature-duration probability in spring

the experimental temperature and duration probability increased, the mortality rate of *N. albiflora* showed an upward trend, with the highest mortality rate observed in the 8.25°C-100% duration probability treatment group. Significant differences in survival rates were found between the 4.0°C-75%, 4.0°C-100% and 8.5°C-75%, 8.5°C-100% elevated temperature-duration probability experimental groups (Figure 5).

Under summer temperature conditions, no mortality was observed in the control group, the various 0.5°C rise duration probability experimental groups, and the 1.0°C-25%

elevated temperature-duration probability experimental group of *N. albiflora*. Mortality occurred in all other experimental groups, with average survival rates for 1.0°C-50%, 1.0°C-75%, 1.0°C-100%, 2.0°C-25%, 2.0°C-50%, 2.0°C-75%, 2.0°C-100%, 4.0°C-25%, 4.0°C-50%, 4.0°C-75%, 4.0°C-100%, 8.5°C-25%, 8.5°C-50%, 8.5°C-75%, and 8.5°C-100% being (90.0±3.3)%, (90.0±5.8)%, (90.0±3.3)%, (90.0±5.8)%, (90.0±3.3)%, (84.4±5.0)%, (84.4±8.8)%, (90.0±6.7)%, (80.0±6.7)%, (70.0±3.3)%, (70.0±3.8)%, (83.3±3.3)%, (70.0±1.9)%, (70.0±3.3)%, and (61.1±6.9)% respectively. Overall, as the experimental temperature and




Fig. 6. Survival rate of *N. albiflora* at different elevated temperature amplitudes and duration probability in summer

duration probability increased, the mortality rate of *N. albiflora* generally showed an upward trend, with the highest mortality rate observed in the 8.25°C-100% duration probability treatment group, where the average mortality rate was 35%. Significant differences in survival rates were found between 2.0°C-75% and 2.0°C-100% compared to 2.0°C-25% and 2.0°C-50%; 4.0°C-75% and 4.0°C-100% compared to 4.0°C-25% and 4.0°C-50%; and 8.5°C-50% and 8.5°C-75% compared to 8.5°C-25% and 8.5°C-100% elevated temperature-duration probability experimental groups (Figure 6).

Under autumn temperature conditions, mortality was observed in the experimental groups with elevated temperature-duration probabilities of 1.0°C-75%, 1.0°C-100%, 2.0°C-75%, 2.0°C-100%, 4.0°C-50%, 4.0°C-75%, 4.0°C-100%, 8.5°C-25%, 8.5°C-50%, 8.5°C-75%, and 8.5°C-100%, with average survival rates of (94.4±3.8)%, (90.0±5.8)%, (93.3±3.8)%, (90.3±6.7)%, (93.3±0.0)%, (84.4±8.8)%, (84.4±6.7)%, (86.7±0.6)%, (83.3±3.3)%, (83.3±6.1)%, and (76.7±8.4)% respectively. No mortality occurred in the control group or the other elevated temperature-duration probability experimental groups of *N. albiflora*. Overall, as the experimental temperature and duration probability increased, the mortality rate of *N. albiflora* showed an upward trend, with the highest mortality rate observed in the 8.5°C-100% duration probability treatment group. Significant differences in survival rates were noted between 1.0°C-75% and 1.0°C-100%; 2.0°C-75% and 2.0°C-100%; 4.0°C-50% and 4.0°C-75%, 4.0°C-100%; and 8.5°C-50%, 8.5°C-75% compared to 8.5°C-25%, 8.5°C-100% elevated temperature-duration probability experimental groups (Figure 7).

Under winter temperature conditions, mortality was observed only in the *N. albiflora* experimental groups with

8.5°C-75% and 8.5°C-100% elevated temperature-duration probabilities, with average survival rates of (94.4±1.9) % and (85.6±6.9) % respectively. No mortality occurred in the control group or the other experimental groups of *N. albiflora*, with the highest mortality rate observed in the 8.5°C-100% treatment group. Significant differences in survival rates were found between the 8.5°C-75% and 8.5°C-100% elevated temperature-duration probability experimental groups (Figure 8).

THE IMPACT OF PROBABILISTIC FREQUENCY OF TEMPERATURE INCREMENT ON THE SURVIVAL RATE OF *T. OVATUS* ACROSS DIFFERENT SEASONS

The two-factor analysis of variance results for the effects of elevated temperature-duration probability on the survival rates of *T. ovatus* and *N. albiflora* (Table 1) revealed that, in spring, elevated temperature had no significant effect on the survival rate of *N. albiflora*, whereas duration probability significantly affected it. There was an interaction between elevated temperature and duration probability. In summer and autumn, both elevated temperature and duration probability significantly influenced the survival rate of *N. albiflora*, with an interaction between these two factors. In winter, neither elevated temperature nor duration probability significantly affected the survival rate of *N. albiflora*, but there was an interaction between them. In summer, elevated temperature significantly affected the survival rate of *T. ovatus*, while duration probability did not have a significant effect, indicating an interaction between elevated temperature and duration probability.




Fig. 7. Survival rate of *N. albiflora* at different elevated temperature amplitudes and duration probability in autumn




Fig. 8. Survival rate of *N. albiflora* at different elevated temperature amplitudes and duration probability in winter

DISCUSSION

Temperature is an extremely important factor in aquatic environments. As one of the primary variables in physiological ecology, temperature is related to the metabolic processes of poikilothermic animals' basic life history. It is one of the key factors controlling poikilotherms' life cycle and behavioral changes through physiological and bio-

chemical means.¹¹ Fish lack the ability to regulate their body temperature, so their body temperature changes with the surrounding temperature, leading to changes in growth, behavior, reproduction, metabolic activities, lifespan, and survival rates.¹² Generally, fish may seek optimal temperature conditions as their habitat, and significant temperature changes under optimal conditions can be considered a stress source for fish behavior.^{13,14} Therefore, thermal

Table 1. Two-way ANOVA results of the effects of elevated temperature amplitudes and duration probability on the survival rate of *T. ovatus* and *N. albiflora*

Species	Season	Source of variations	df	MS	F	P
<i>N. albiflora</i>	Spring	Elevated temperature (°C)	4	0.31	2.77	0.08
		Duration probability (°C)	3	0.64	5.71	0.01
		Interaction of elevated temperature×duration probability	12	0.11	144.18	0.00
	Summer	Elevated temperature (°C)	4	4.8	23.44	0.00
		Duration probability (°C)	3	1.22	5.93	0.01
		Interaction of elevated temperature×duration probability	12	0.21	90.00	0.00
	Autumn	Elevated temperature (°C)	4	571.29	18.91	0.00
		Duration probability (°C)	3	257.83	8.54	0.00
		Interaction of elevated temperature×duration probability	12	30.21	111.12	0.00
	Winter	Elevated temperature (°C)	4	0.17	2.18	0.13
		Duration probability (°C)	3	0.08	1.00	0.43
		Interaction of elevated temperature×duration probability	12	0.08	169.18	0.00
<i>T. ovatus</i>	Summer	Elevated temperature (°C)	4	0.2868	5.6984	0.01
		Duration probability (°C)	3	0.05	1.00	0.43
		Interaction of elevated temperature×duration probability	12	0.05	179.16	0.00

emissions from power plants, including nuclear power plants, directly impact fish.

The findings of this study reveal that in different seasons, the mortality rates of *T. ovatus* and *N. albiflora* both escalated with increasing acclimation temperatures. When the temperature increase amplitude was small, the mortality rate was low or there was no mortality; however, as the temperature increase became more significant, the mortality rate noticeably increased. This is related to the temperature tolerance of fish, as different species have varying upper and lower limits of temperature tolerance. Within the optimal temperature range, a moderate temperature increase can enhance the feeding ability of marine fish, promote their sexual maturity, and accelerate growth. However, once the temperature exceeds their tolerance range, it may culminate in mortality.^{15,16} In this experiment, within the same season, the survival rate of *T. ovatus* under the same temperature increase amplitudes and duration probabilities (Figures 1-8) was significantly higher than that of *N. albiflora*. This is because the lethal temperatures, critical temperatures, and thermal adaptation capacities differ among fish species, closely related to the latitude, geographic distribution, and the temperature of their birth habitats. Generally, warm-water fish have a greater heat tolerance than cold-water species.¹⁷ The *T. ovatus* is a more typical southern fish species,⁹ while the *N. albiflora* is more typical of northern regions.¹⁰ The habitat temperature of the *T. ovatus* in different seasons is significantly higher than that of the *N. albiflora*, making the *N. albiflora* more sensitive to temperature increment compared to the *T. ovatus*. Specifically, the suitable temperature range for *T. ovatus* is about 25-32°C,¹⁸ while for *N. albiflora*, it is approx-

imately 20-25°C.¹⁹ Therefore, in spring, autumn, and winter, the acclimation temperatures combined with different temperature increment still fall within the suitable temperature range for *T. ovatus*. Only in summer, when the acclimation temperature is combined with an 8.5°C temperature increase, do some experimental groups experience lower mortality. The suitable temperature range for *N. albiflora* is relatively narrow; in summer and autumn, compared to *T. ovatus*, even a lower temperature increase combined with the acclimation temperature exceeds the suitable temperature range for *N. albiflora*, leading to mortality. However, in spring and winter, the water temperature, combined with different temperature increment for *N. albiflora*, generally remains below its suitable temperature range. Given that the experimental methodology of this study differs from typical fish tolerance tests—placing fish under thermal shock at predetermined experimental temperatures—the main cause of mortality is likely the stress response caused by sudden thermal shock under low-temperature conditions.

Fish can perceive changes in water temperature and regulate their body temperature within a narrow adaptive range through physiological and ecological behaviors to avoid harm from extreme high temperatures.²⁰ Whether exposure to temperatures near their lethal limits results in death depends on the duration of exposure to such temperatures, a phenomenon that is more evident in the experiments with *N. albiflora*. The results of this study show that at lower temperature increase amplitudes, the duration of elevated temperature has a minimal impact on the survival rate of *N. albiflora*. However, as the temperature increase amplitude grows, the effect of elevated temperature

duration on the survival rate of *N. albiflora* becomes more significant, indicating that the longer the duration of elevated temperature, the lower the survival rate. Increased temperatures can mobilize the rates of metabolic enzymes, ion channels, mitochondria, and other important biochemical reaction processes in the organism to compensate for the impact of temperature stress. However, prolonged exposure beyond the organism's maximum physiological adjustment limit leads to a weakening or collapse of physiological compensation mechanisms, ultimately resulting in death.²¹

Previous studies assessing the impact of thermal power plant thermal discharge on aquatic organisms, such as fish, mainly focused on the tolerance of aquatic life under different temperature increase amplitudes, often overlooking the interaction between actual operating conditions and hydrological conditions over time. The results of this study indicate that in assessing the impact of thermal power plants on fish, especially for temperature-sensitive species, to enhance the accuracy of the assessment, it is not only necessary to consider the temperature increase amplitudes but also the duration of different temperature increase amplitudes as an important factor to be taken into account.

CONCLUSION

The mortality patterns of *T. ovatus* and *N. albiflora* vary across seasons and in response to temperature changes. *T. ovatus* shows remarkable resilience, with no mortality in spring, autumn, and winter, and only succumbing in summer. In contrast, *N. albiflora* experiences mortality throughout the year, with the most severe losses also in summer. Both species exhibit higher mortality rates with increasing temperatures and longer exposure times, indicating that the survival rates of these fish are influenced by both the magnitude and the duration of temperature increment. *T. ovatus* demonstrates a higher tolerance to temperature

fluctuations than *N. albiflora*, suggesting a species-specific capacity to withstand environmental changes.

AUTHORS' CONTRIBUTION

Conceptualization: Lei Li (Lead). Investigation: Lei Li (Lead). Project administration: Chenshan Shao (Lead). Supervision: Jiaying Cai (Lead). Validation: Jiaying Cai (Lead). Formal Analysis: Baojun Tang. Visualization: Baojun Tang (Lead). Writing – original draft: Baojun Tang (Lead). Data curation: Weiwei Su (Lead). Resources: Weiwei Su (Lead). Software: Weiyi Zou (Lead). Methodology: Weiyi Zou (Lead). Writing – review & editing: Mei Jiang (Lead).

ETHICAL CONDUCT APPROVAL – IACUC

RESEARCH INVOLVING ANIMALS

Our experiments involving vertebrates were conducted within the ethical guidelines provided by our institution and national or international regulations. We confirm that all efforts were made to ameliorate animals' suffering during the experiments.

INFORMED CONSENT STATEMENT

All authors and institutions have confirmed this manuscript for publication.

DATA AVAILABILITY STATEMENT

All are available upon reasonable request.

Submitted: April 19, 2024 CDT, Accepted: May 28, 2024 CDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-SA-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc-sa/4.0> and legal code at <https://creativecommons.org/licenses/by-nc-sa/4.0/legalcode> for more information.

REFERENCES

1. Argüelles R, Toledo M, Martín MA. Study of the Tagus River and Entrepeñas reservoir ecosystem around the Trillo nuclear power plant using chemometric analysis: Influence on water, sediments, algae and fish. *Chemosphere*. 2021;279:130532. [doi:10.1016/j.chemosphere.2021.130532](https://doi.org/10.1016/j.chemosphere.2021.130532)
2. Mari C. The costs of generating electricity and the competitiveness of nuclear power. *Progress in Nuclear Energy*. 2014;73:153-161. [doi:10.1016/j.pnucene.2014.02.005](https://doi.org/10.1016/j.pnucene.2014.02.005)
3. Macknick J, Newmark R, Heath G, Hallett KC. Operational water consumption and withdrawal factors for electricity generating technologies: a review of existing literature. *Environmental Research Letters*. 2012;7(4):045802. [doi:10.1088/1748-9326/7/4/045802](https://doi.org/10.1088/1748-9326/7/4/045802)
4. Teixeira TP, Neves LM, Araújo FG. Effects of a nuclear power plant thermal discharge on habitat complexity and fish community structure in Ilha Grande Bay, Brazil. *Marine Environmental Research*. 2009;68(4):188-195. [doi:10.1016/j.marenvres.2009.06.004](https://doi.org/10.1016/j.marenvres.2009.06.004)
5. Chen XJ, Liu JL, Lin DM, Gong Y. Review of development and trend in fisheries resource science. *Journal of Shanghai Ocean University*. 2012;31(5):1168-1179. [doi:10.12024/jsou.20220503856](https://doi.org/10.12024/jsou.20220503856)
6. Payne NL, Smith JA, van der Meulen DE, et al. Temperature dependence of fish performance in the wild: links with species biogeography and physiological thermal tolerance. *Functional Ecology*. 2016;30(6):903-912. [doi:10.1111/1365-2435.12618](https://doi.org/10.1111/1365-2435.12618)
7. Dent L, Lutterschmidt WI. Comparative thermal physiology of two sympatric sunfishes (Centrarchidae: Perciformes) with a discussion of microhabitat utilization. *Journal of Thermal Biology*. 2003;28(1):67-74. [doi:10.1016/S0306-4565\(02\)00038-4](https://doi.org/10.1016/S0306-4565(02)00038-4)
8. Pang X, Cao ZD, Peng JL, Fu SJ. The effects of feeding on the swimming performance and metabolic response of juvenile southern catfish, *Silurus meridionalis*, acclimated at different temperatures. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2010;155(2):253-258. [doi:10.1016/j.cbpa.2009.11.005](https://doi.org/10.1016/j.cbpa.2009.11.005)
9. Liu B, Guo HY, Zhu KC, et al. Growth, physiological, and molecular responses of golden pompano *Trachinotus ovatus* (Linnaeus, 1758) reared at different salinities. *Fish Physiology and Biochemistry*. 2019;45(6):1879-1893. [doi:10.1007/s10695-019-00684-9](https://doi.org/10.1007/s10695-019-00684-9)
10. Han ZQ, Gao TX, Yanagimoto T, Sakurai Y. Genetic population structure of *Nibea albiflora* in Yellow Sea and East China Sea. *Fisheries Science*. 2008;74(3):544-552. [doi:10.1111/j.1444-2906.2008.01557.x](https://doi.org/10.1111/j.1444-2906.2008.01557.x)
11. Poole GC, Berman CH. An ecological perspective on in-stream temperature: natural heat dynamics and mechanisms of human-caused thermal degradation. *Environmental Management*. 2001;27(6):787-802. [doi:10.1007/s002670010188](https://doi.org/10.1007/s002670010188)
12. Kumar N, Thorat ST, Gite A, Patole PB. Selenium nanoparticles and omega-3 fatty acid enhanced thermal tolerance in fish against arsenic and high temperature. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 2022;261:109447. [doi:10.1016/j.cbpc.2022.109447](https://doi.org/10.1016/j.cbpc.2022.109447)
13. Logan LH, Stillwell AS. Water temperature duration curves for thermoelectric power plant mixing zone analysis. *Journal of Water Resources Planning and Management*. 2018;144(9):04018058. [doi:10.1061/\(ASCE\)WR.1943-5452.0000980](https://doi.org/10.1061/(ASCE)WR.1943-5452.0000980)
14. Li S, Guo H, Chen Z, et al. Effects of acclimation temperature regime on the thermal tolerance, growth performance and gene expression of a cold-water fish, *Schizothorax prenanti*. *Journal of Thermal Biology*. 2021;98:102918. [doi:10.1016/j.jtherbio.2021.102918](https://doi.org/10.1016/j.jtherbio.2021.102918)
15. Kir M, Sunar MC, Altındağ BC. Thermal tolerance and preferred temperature range of juvenile meagre acclimated to four temperatures. *Journal of Thermal Biology*. 2017;65:125-129. [doi:10.1016/j.jtherbio.2017.02.018](https://doi.org/10.1016/j.jtherbio.2017.02.018)
16. Rajaguru S, Ramachandran S. Temperature tolerance of some estuarine fishes. *Journal of Thermal Biology*. 2001;26(1):41-45. [doi:10.1016/S0306-4565\(00\)00024-3](https://doi.org/10.1016/S0306-4565(00)00024-3)
17. Payne NL, Morley SA, Halsey LG, et al. Fish heating tolerance scales similarly across individual physiology and populations. *Communications Biology*. 2021;4(1):264. [doi:10.1038/s42003-021-01773-3](https://doi.org/10.1038/s42003-021-01773-3)

18. Tan X, Lin H, Huang Z, et al. Effects of dietary leucine on growth performance, feed utilization, non-specific immune responses and gut morphology of juvenile golden pompano *Trachinotus ovatus*. *Aquaculture*. 2016;465:100-107. [doi:10.1016/j.aquaculture.2016.08.034](https://doi.org/10.1016/j.aquaculture.2016.08.034)

19. Nóbrega MF, Oliveira MA, Lira MG, Rocha SdS, Lins Oliveira JE. Sustainability of tunas and swordfish exploitation in the equatorial tropical Atlantic Ocean. *Marine Policy*. 2023;155:105755. [doi:10.1016/j.marpol.2023.105755](https://doi.org/10.1016/j.marpol.2023.105755)

20. Reyes I, Díaz F, Re AD, Pérez J. Behavioral thermoregulation, temperature tolerance and oxygen consumption in the Mexican bullseye puffer fish, *Sphoeroides annulatus* Jenyns (1842), acclimated to different temperatures. *Journal of Thermal Biology*. 2011;36(3):200-205. [doi:10.1016/j.jtherbio.2011.03.003](https://doi.org/10.1016/j.jtherbio.2011.03.003)

21. Matoo OB, Lannig G, Bock C, Sokolova IM. Temperature but not ocean acidification affects energy metabolism and enzyme activities in the blue mussel, *Mytilus edulis*. *Ecology and Evolution*. 2021;11(7):3366-3379. [doi:10.1002/ece3.7289](https://doi.org/10.1002/ece3.7289)