

The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

Editor-in-Chief

Dan Mires

Editorial Board

Sheenan Harpaz Agricultural Research Organization
Beit Dagan, Israel

Zvi Yaron Dept. of Zoology
Tel Aviv University
Tel Aviv, Israel

Angelo Colorni National Center for Mariculture, IOLR
Eilat, Israel

Rina Chakrabarti Aqua Research Lab
Dept. of Zoology
University of Delhi

Ingrid Lupatsch Swansea University
Singleton Park, Swansea, UK

Jaap van Rijn The Hebrew University
Faculty of Agriculture
Israel

Spencer Malecha Dept. of Human Nutrition, Food
and Animal Sciences
University of Hawaii

Daniel Golani The Hebrew University of Jerusalem
Jerusalem, Israel

Emilio Tibaldi Udine University
Udine, Italy

Copy Editor

Ellen Rosenberg

Published under auspices of
**The Society of Israeli Aquaculture and
Marine Biotechnology (SIAMB),
University of Hawaii at Manoa Library**

and
**University of Hawaii Aquaculture
Program** in association with
AquacultureHub

<http://www.aquaculturehub.org>



UNIVERSITY
of HAWAII
MĀNOA
LIBRARY



AquacultureHub
educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:
Israeli Journal of Aquaculture - BAMIGDEH -
Kibbutz Ein Hamifratz, Mobile Post 25210,
ISRAEL

Phone: + 972 52 3965809

<http://siamb.org.il>

Isolation of Putative Probiotics from the Intestines of Indian Major Carps

Shubhadeep Ghosh*, Archana Sinha and Chittaranjan Sahu**

Central Institute of Fisheries Education, Kolkata Center 32, GN Block, Sector V,
Salt Lake City, Kolkata 700091, India

(Received 25.11.06, Accepted 23.2.07)

Key words: probiotics, antibacterial activity, *Bacillus subtilis*, non-pathogen, isolation, adherence, biochemical characterization

Abstract

Probiotic strains were isolated from the intestines of 28 Indian major carps (*Labeo rohita*, *Catla catla*, *Cirrhinus mrigala*) cultured in sewage-fed ponds. Intestinal microflora were isolated, counted, and identified, and their *in vitro* antibacterial properties were determined. The double layer method revealed that four strains (SG1, SG2, SG3, SG4) elicited antagonism against the pathogen *Pseudomonas fluorescens*. Using cross streak and parallel streak methods, SG4 maximally inhibited the fish pathogens *Pseudomonas fluorescens*, *Aeromonas hydrophila*, and *Edwardsiella tarda*. Therefore, SG4 was chosen for further study as a probiotic. SG4 was non-pathogenic to *C. mrigala* fingerlings, and adhered well to stainless steel, high density polyethylene, and glass plate substrates. Biochemical tests showed that SG4 was endospore-forming, aerobic, motile, rod-shaped *Bacillus subtilis*. However, since *in vitro* experiments can only give an indication of *in vivo* success, future research should include large-scale *in vivo* experiments.

Introduction

Disease is a major problem in the fish farming industry. Although vaccines are being developed and marketed, they cannot be used as a universal disease control measure in aquaculture. The use of antibiotics to cure bacterial infection and prevent fish mortality in aquaculture is becoming limited as pathogens develop resistance to the drugs (Alderman and Hastings, 1998). Further, beneficial bacterial flora are killed or inhibited by orally adminis-

tered antibiotics, leading to efforts to find alternative disease prevention methods such as the use of non-pathogenic bacteria as probiotic biocontrol agents.

Verschuere et al. (2000) defined probiotics as "a live microbial adjunct which has a beneficial effect on the host by modifying the host-associated or ambient microbial community, by ensuring improved use of the feed or enhancing its nutritional value, by enhancing the host

* Corresponding author. Veraval Regional Center of CMFRI, Bhidiya, Veraval, Gujarat 362269, India. E-mail: shubhadeep_1977@yahoo.com; shubha4444@gmail.com

** Current address: Department of Zoology, University of Kalyani, Nadia, West Bengal 741235, India

response towards disease, or by improving the quality of its ambient environment”.

The use of commercial probiotics in fish is relatively ineffective as most commercial preparations are based on strains isolated from non-fish sources that are unable to survive or remain viable at high cell density in the intestinal environment of fish during the active growth phase of the fish (Moriarty, 1996). Hence, there is elegant logic in isolating putative probiotics from the host in which the probiotic is intended for use. Such strains should perform better because they have already adhered to the gut wall of the fish and, thus, are well-adapted to compete with pathogens for nutrients. Presumably, strains that develop dominant colonies in the fish intestine are good candidates for preventing the adhesion of pathogens on the gut wall. The present study was undertaken to isolate probiotic bacterial strains from fish intestines and screen them by *in vitro* testing of their antagonism to pathogens and attachment to substrates.

Materials and Methods

Isolation of bacterial strains. Fingerlings and juveniles of Indian major carps, *Labeo rohita*, *Catla catla*, and *Cirrhinus mrigala*, were collected at regular intervals from the Nalban sewage-fed fisheries at Salt Lake City, Kolkata. They were brought to the laboratory alive and sacrificed. The ventral surfaces were sterilized using 70% ethanol and aseptically dissected to remove the intestines. The intestines were opened by a longitudinal incision and thoroughly flushed with sterilized chilled normal saline solution (NSS) to remove feed materials, dirt, and other impurities. Excess moisture was blotted with filter paper and the intestines were weighed, macerated with a sterile glass rod, and homogenized in sterile NSS (1:10; wt:vol) using a vortex mixer. Samples of the thoroughly macerated and homogenized intestines were serially diluted in NSS and aseptically plated by the spread plate technique on All Purpose Tween's Agar (APTA; Evans and Niven, 1951) to isolate putative probiotic bacteria and on Nutrient Agar (NA; Hi-Media) to determine total plate count. The inoculated agar plates

were aerobically incubated at 35°C for 48 h and the total bacteria were counted.

Assay for antibacterial activity. The double layer method reported by Dopazo et al. (1988) was used as an inhibition test to identify potential probiotics against the fish pathogen *Pseudomonas fluorescens*. Bacterial strains that produced a clear inhibitory zone with a diameter at least 5 mm greater than that of the *P. fluorescens* colony were further examined for antagonistic properties against three fish pathogens, *P. fluorescens*, *Aeromonas hydrophila*, and *Edwardsiella tarda* by cross streak assay (Austin et al., 1992) and the parallel streak method (Nakamura et al., 1999). The pathogens were provided by the Central Institute of Fisheries Education in Mumbai, India. Their identities were confirmed following the procedures of Krieg and Holt (1984).

Determination of pathogenicity of the most effective probiotic strain. The pathogenicity of the most effective probiotic strain was determined by challenging *C. mrigala* fingerlings (avg wt 6.6±1.3 g, avg length 7.8±1.4 cm) in an immersion assay and observing their health status (Austin et al., 1995). Fingerlings were procured from commercial fish breeders in Kolkata and acclimatized in laboratory conditions for five days. The experiment was carried out in triplicate 500-l circular FRP tanks for ten days. Fingerlings were stocked at 30 per tank and maintained in uniform water conditions with aeration and filtration. The probiotic strain was grown for 24 h in tryptone soya broth (TSB; Hi-Media). Cells were harvested by centrifugation at 6000 rpm (2000 x g) for 20 min at 4°C in a cooling centrifuge (C-23, Remi, Mumbai, India). The cells were washed and resuspended in sterile saline. The number of cells was estimated by spread plating on tryptone soya agar (TSA; Hi-Media) after incubation at 35°C for 48 h. The remaining probiotic cells in the sterile saline were added to the fish tanks to a level of 10⁷ cells/ml. Fish in control tanks received a sterile saline solution lacking bacterial inoculum. All fish were fed a basal diet (35% protein) at 5% of their body weight in two doses. At the end of the experiment, gross pathological features, abnormalities, and mortality rates, considered to be due to the inoculation, were recorded.

Adherence or biofilm formation of the most effective probiotic strain on substrates. The ability of the most effective of the isolated probiotics to adhere to and/or form a biofilm on stainless steel, high density polyethylene (HDPE), and glass plates was tested as described by Karunasagar et al. (1996) and Venugopal et al. (1999).

Biochemical characterization and identification of the most effective probiotic strain. The most effective probiotic was characterized and identified based on Gram reaction; spore formation; cellular morphology; motility; growth in different salinities and pH; sugar utilization; amino acid decarboxylation; catalase and oxidase production; nitrate reduction; hydrogen sulfide production; starch, casein, and urea hydrolysis; gelatin liquefaction; and IMVIC tests. The biochemical tests were performed according to Collins et al. (1989) and Pacarynuk and Danyk (2004).

Results

Isolation of bacterial strains. The intestines of 28 fish were screened to isolate probiotic bacteria. The sampled fish ranged 11-27 cm (avg 17.5±4.8 cm) and 12-150 g (avg 57.88±49.03 g). The log₁₀ values of total plate count (TPC) per gram intestine ranged 7.544-9.531 on NA (avg 8.605±0.525) and 5.114-6.978 on APTA (avg 6.019±0.602).

Assay for antibacterial activity. The double layer method showed that four strains, SG1 isolated from *C. mrigala* (23 cm; 100 g), SG2 isolated from *C. catla* (18 cm; 60 g), SG3 isolated from *C. catla* (27 cm; 140 g), and SG4 isolated from *C. mrigala* (23 cm; 115 g) produced a zone of inhibition with a diameter that was more than 5 mm greater than the diameter of the *P. fluorescens* colony.

The cross streak method revealed that the two strains from *C. catla* (SG2 and SG3) did not inhibit the pathogenic test strains. The two strains from *C. mrigala* (SG1 and SG4) had inhibition zones of 5.21±0.57 mm and 5.90±3.37 mm against *P. fluorescens*, 4.05±1.73 mm and 3.16±0.64 mm against *A. hydrophila*, and 3.48±1.16 mm and 3.8±1.27 mm against *E. tarda*, respectively. The antibacterial activity of SG1 and SG4 was further confirmed

using the parallel streak method against the same three pathogenic test strains. For SG1, there was no noticeable reduction in the width of any pathogen streak on the test plates compared to that of the control. SG4, however, markedly reduced the width of all pathogen streaks when compared to the control.

By virtue of its greater inhibition zone in the cross streak method and positive antibacterial activity in the parallel streak method, SG4 was further tested in the following experiments.

Determination of pathogenicity of SG4. The addition of SG4 cells to the rearing water of *C. mrigala* fingerlings produced no gross pathological symptoms, abnormalities, or mortality, confirming its non-pathogenicity.

Adherence or biofilm formation of SG4 on substrates. The adherence or biofilm formation of SG4 was highest on HDPE, followed by stainless steel and glass. The log₁₀ counts on HDPE, steel, and glass were 3.316/cm², 2.592/cm², and 0.47/cm², respectively.

Biochemical characterization and identification of SG4. SG4 had a strong gram-positive reaction. Cellular morphology suggests that it is a thick unicellular rod-shaped bacteria, capable of forming round, wavy, convex, rough, and opaque colonies. Cells are aerobic, endospore-forming, and motile while spores are ellipsoidal and central in position. SG4 is capable of utilizing glucose, xylose, sucrose, mannitol, arabinose, and fructose as carbohydrate sources for growth and grew at pH levels of 5, 8, 9, and 11 and salinities of 3%, 6%, and 8%, but not 10%. It did not decarboxylate the amino acids lysine, ornithine, and arginine. SG4 was positive for the Vogues Proskauer reaction and negative for the methyl red test. It can produce catalase, oxidase, and indole, but not hydrogen sulfide and can utilize citrate, reduce nitrate, liquefy gelatin, and hydrolyze starch and casein, but not urea.

Discussion

Isolation of bacterial strains. The mean log₁₀ intestinal count on NA, 8.605±0.525/g, agrees with that of Galindo (2004), who reported a mean log₁₀ intestinal count of 8.09±0.43/g in 17 species of carnivorous, herbivorous, and

omnivorous freshwater fishes, and Ringo et al. (1995), who found that the digestive tract of fin-fish contains 10^8 cells/g, equivalent to 8/g when converted into a logarithmic scale with a base of 10. The mean \log_{10} count of lactic acid bacteria on APTA, $6.019 \pm 0.602/g$, was higher than that obtained by Galindo (2004) who obtained a mean \log_{10} intestinal lactic acid bacteria (LAB) count of $4.72 \pm 1.4/g$ while working on 17 freshwater fish species. The fish intestine is a favorable ecological niche for microorganisms, which reach much higher numbers than in the surrounding water (Austin and Austin, 1987).

Assay for antibacterial activity. The double layer method showed that only four strains of all those isolated from the intestines of 28 fish had inhibition zones that exceeded the circumference of a colony of pathogenic *P. fluorescens* by at least 5 mm. A similar frequency of inhibitory bacteria was observed for isolates from halibut larvae (Bergh, 1995), rainbow trout (Brunt and Austin, 2005), turbot (Hjelm et al., 2004), and shrimp (Rengpipat et al., 1998). The inhibition was caused by the release of chemical substances with bactericidal or bacteriostatic effects on *P. fluorescens*.

The inhibition zones exhibited by SG1 and SG4 in the cross streak method were similar to those reported by Abraham (2004) for *Alteromonas* sp. P7 (4.0 ± 0.25 mm) and by Rengpipat et al. (1998) for *Bacillus* strain BY-9 (3-6 mm) against the pathogenic *Vibrio harveyi*. In another study, Galindo (2004) reported *Lactobacillus plantarum* 44a exhibiting inhibition zones of 12.4 ± 1.3 mm and 7.3 ± 0.90 mm against *A. hydrophila* at pH levels of 3.7 and 4.5, respectively, and *Lactobacillus lactis* 18f exhibiting inhibition zones of 11.2 ± 1.16 mm, 2.0 mm, and 8.9 ± 2.1 mm against *A. hydrophila*, *E. tarda*, and *Staphylococcus aureus*, respectively, at pH 6.0. The inhibition seen in the cross streak method resulted from interruption of pathogen growth caused by production of one or more of the following factors: antibiotics, antimicrobial peptides, bacteriocins, siderophores, lysosomes, proteases, hydrogen peroxide, and organic acids (Verschuere et al., 2000).

SG4, but not SG1, reduced the width of pathogen streaks, in agreement with Nakamura et al. (1999) who reported that only twelve of 51

bacterial isolates with suppressive activity had inhibitory effects on pathogen growth in the parallel streak method. The reduction of pathogen growth and cell density indicate that extracellular bacteriolytic products produced by the SG4 were responsible for this inhibition.

The *in vitro* production of compounds that inhibit known pathogens is often used in the selection of putative probiotic strains (Rengpipat et al., 1998). In the present *in vitro* study, SG4 inhibited all three pathogenic strains and hence was selected for further study as probiont.

Determination of pathogenicity of SG4. There is no evidence that SG4 has any harmful effect on *C. mrigala* fingerlings since there was no mortality and a complete absence of any pathology or abnormality in the fingerlings. Austin et al. (1995) also found that isolated probiotic strains were non-pathogenic. The non-pathogenicity of SG4 ensures that it is safe for use as a probiotic.

Adherence or biofilm formation of SG4 on substrates. The adherence or biofilm formation of SG4 on three substrates was similar to the findings of Faille et al. (2002), who observed a higher number of adhered cells on the hydrophobic materials HDPE and Teflon, but not much difference (less than 1 log unit) between these two materials. The ability to adhere to a substrate is an important criterion for selection of a probiont because adhesion to host surfaces is essential for effective colonization of the probiont in the host and interruption of adhesion of pathogens is essential for bio-control of the pathogen (Chabrillon et al., 2005).

Biochemical characterization and identification of SG4. The morphological and physiological characteristics of SG4 and comparison with Sneath (1986) and Priest (1993) place this bacterium in the *Bacillus* genus and species *subtilis*. *Bacillus* is distinguished from other endospore-forming bacteria by being a strict or facultative aerobe, rod-shaped, and catalase-positive. *Bacillus subtilis* is distinguished from other members of the genus by its biochemical reactions, unicellular rod-shaped morphology, and ellipsoidal spore-forming ability.

Bacilli are not autochthonous to the gas-

trointestinal tract but have been isolated from carps (Kumar et al., 2006), coastal fishes (Sugita et al., 1998), shrimps (Sharmila et al., 1996), bivalves (Sugita et al., 1981), shrimp culture ponds (Vaseeharan and Ramasamy, 2003), and shrimp larvae-rearing medium (Rengpipat et al., 1998). Most *Bacilli* produce antibiotics such as difficidin, oxydifficidin, bacitracin, polymyxin, subtilin, mycobacillin, bacillin, gramicidin, or bacillomycin B and are antagonistic to pathogenic bacteria in both *in vivo* and *in vitro* conditions (Korzybski et al., 1978; Zimmerman et al., 1987). *Bacilli* are not associated with aquatic organism pathology and are widely accepted and used as probiotics (Moriarty, 1998).

In this study we isolated probiotics from fish intestines and screened their potential to combat fish pathogens in aquaculture in *in vitro* experiments. The use of a probiotic originating from the fish intestine will enable the probiotic to proliferate rapidly once it reaches the fish intestine as a feed supplement. While *in vitro* experiments can indicate success *in vivo*, assumptions based on *in vitro* experiments may not be true for *in vivo* modes of action. Therefore, future research should include large scale *in vivo* experiments along with study of the mechanism of antagonistic action between the probiotic and the pathogen.

Acknowledgement

The authors express their gratitude to the Indian Council of Agricultural Research, New Delhi, for financing this study under an AP-Cess Fund Project.

References

- Abraham T.J.**, 2004. Antibacterial marine bacterium deter luminous Vibriosis in shrimp larvae. *NAGA* 27(3&4): 28-31.
- Alderman D.J. and T.S. Hastings**, 1998. Antibiotic use in aquaculture: development of antibiotic resistance-potential for consumer health risks. *Int. J. Food Sci. Technol.*, 33:139-155.
- Austin B. and D.A. Austin**, 1987. *Bacterial Fish Pathogens: Diseases in Farmed and Wild Fish*. Ellis-Horwood Ltd., Chichester, UK. pp. 112-117.
- Austin B., Baudet E. and M. Stobie**, 1992. Inhibition of bacterial fish pathogens by *Tetraselmis suvicica*. *J. Fish Dis.*, 15:55-61.
- Austin B., Stuckey L.F., Robertson P.A.W., Effendi I. and D.R.W. Griffith**, 1995. A probiotic strain of *Vibrio alginolyticus* effective in reducing diseases caused by *Aeromonas salmonicida*, *Vibrio anguillarum* and *Vibrio ordalii*. *J. Fish Dis.*, 18:93-96.
- Bergh O.**, 1995. Bacteria associated with early life stages of halibut, *Hippoglossus hippoglossus* L., inhibit growth of a pathogenic *Vibrio* sp. *J. Fish Dis.*, 18:93-96.
- Brunt J. and B. Austin**, 2005. Use of a probiotic to control lactococcosis and streptococcosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *J. Fish Dis.*, 28:693-701.
- Chabrillon M., Rico R., Balebona M.C. and M.A. Morinigo**, 2005. Adhesion to sole (*Solea senegalensis*) mucus of microorganisms isolated from farmed fish, and their interaction with *Photobacterium damsela* subsp *piscicida*. *J. Fish Dis.*, 28:229-237.
- Collins C.H., Lyne P.M. and J.M. Grange**, 1989. *Microbiological Methods*. 6th ed., Butterworth, London, UK. 409 pp.
- Dopazo C.P., Lemos M.L., Lodeiros C., Bolinches J., Barja J.L. and A.E. Toranzo**, 1988. Inhibitory activity of antibiotic-producing marine bacteria against fish pathogens. *J. Appl. Microbiol.*, 65:97-101.
- Evans J.B. and C.F. Niven**, 1951. Nutrition of the heterofermentative *Lactobacilli* that cause greening of cured meat products. *J. Bacteriol.*, 62:599-603.
- Faille C., Jullien C., Fontaine F., Bellon-Fontaine M.N., Slomianny C. and T. Benezech**, 2002. Adhesion of *Bacillus* spores and *Escherichia coli* cells to inert surfaces: role of surface hydrophobicity. *Can. J. Microbiol.*, 48(8):728-738.
- Galindo A.B.**, 2004. *Lactobacillus plantarum* 44A as a Live Feed Supplement for Freshwater Fish. Ph.D. diss., Wageningen Univ., Wageningen, Netherlands. (www.gcw.nl/dissertations/content.cfm)
- Hjelm M., Bergh O., Riaza A., Nielsen J., Melchiorson J., Jensen S., Duncan H., Ahrens P., Birkbeck H. and L. Gram**, 2004. Selection and identification of autochthonous

- potential probiotic bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. *Syst. Appl. Microbiol.*, 27:360-371.
- Karunasagar I., Otta S.K. and I. Karunasagar**, 1996. Biofilm formation by *Vibrio harveyi* on surfaces. *Aquaculture*, 140:241-245.
- Korzybski T., Kowszyk-Gindifer Z. and W. Kurylowicz**, 1978. Antibiotics isolated from the genus *Bacillus* (Bacillaceae). pp. 1529-1661. In: *Antibiotics - Origin, Nature and Properties*. Vol. III, Am. Soc. Microbiol., Washington DC, USA.
- Krieg N.R. and J.G. Holt**, 1984. *Bergey's Manual of Systematic Bacteriology*. Vol. 1, Williams and Wilkin, Baltimore, MD, USA. pp. 964.
- Kumar R., Mukherjee S.C., Prasad K.P. and A.K. Pal**, 2006. Evaluation of *Bacillus subtilis* as a probiotic to Indian major carp *Labeo rohita* (Ham.). *Aquac. Res.*, 37(12):1215-1221.
- Moriarty D.J.W.**, 1996. Microbial biotechnology: a key ingredient for sustainable aquaculture. *Infofish Int.*, 4/96:29-33.
- Moriarty D.J.W.**, 1998. Control of luminous *Vibrio* species in penaeid aquaculture ponds. *Aquaculture*, 164:351-358.
- Nakamura A., Takahashi K.G. and K. Mori**, 1999. Vibriostatic bacteria isolated from rearing seawater of oyster brood stock: potentiality as biocontrol agents for *Vibriosis* in oyster larvae. *Fish Path.*, 34(3):139-144.
- Pacarynuk L.A. and H.C. Danyk**, 2004. Biochemical Tests. pp. 28-34. In: *Principles of Microbiology (Laboratory Manual)*. Spring, TX, USA
- Priest F.G.**, 1993. Systematics and Ecology of *Bacillus*. pp. 3-16. In: A.L. Sonensheine, J.A. Koch, R. Losick (eds.). *Bacillus subtilis and Other Gram-positive Bacteria: Biochemistry, Physiology and Molecular Genetics*. Am. Soc. Microbiol., Washington DC, USA.
- Rengpipat S., Rukpratanporn S., Piyatiratitivorakul S. and P. Menasveta**, 1998. Probiotics in aquaculture: a case study of probiotics for larvae of the black tiger shrimp (*Penaeus monodon*). pp. 177-181. In: T.W. Flegel (ed.). *Advances in Shrimp Biotechnology*. Natl. Center for Genetic Engineering and Biotechnology, Bangkok.
- Ringo E., Strom E. and J.A. Tabachek**, 1995. Intestinal microflora of salmonids: A Review. *Aquac. Res.*, 26:773-789.
- Sharmila R., Abraham T.J. and V. Sundararaj**, 1996. Bacterial flora of semi-intensive pond-reared *Penaeus indicus* (H. Milne Edwards) and the environment. *J. Aquac. Trop.* 11:193-203.
- Sneath P.H.A.**, 1986. Endospore-forming gram-positive rods and cocci. pp. 1104-1105. In: P.H.A. Sneath, N.S. Mair, M.E. Sharp, J.G. Holt (eds.). *Bergey's Manual of Systemic Bacteriology*. Vol. 2, Williams and Wilkins, Baltimore, MD, USA.
- Sugita H., Tanaami H., Kobashi T. and Y. Deguchi**, 1981. Bacterial flora of coastal bivalves. *Bull. Jpn. Soc. Sci. Fish.*, 47:655-661.
- Sugita H., Hirose Y., Matsuo N. and Y. Deguchi**, 1998. Production of the antibacterial substance by *Bacillus* sp. strain NM 12, an intestinal bacterium of Japanese coastal fish. *Aquaculture*, 165(3-4):269-280.
- Vaseeharan B. and P. Ramasamy**, 2003. Control of pathogenic *Vibrio* spp. by *Bacillus subtilis* BT23, a possible probiotic treatment for black tiger shrimp *Penaeus monodon*. *Letters Appl. Microbiol.*, 36(2):83-87.
- Venugopal M.N., Karunasagar I., Karunasagar I. and M.C. Varadaraj**, 1999. Effect of sanitizers on *Vibrio parahaemolyticus* in biofilm on stainless steel surface. *Indian J. Microbiol.*, 39:253-254.
- Verschuere L., Rombaut G., Sorgeloos P. and W. Verstraete**, 2000. Probiotic bacteria as biocontrol agents in aquaculture. *Microbiol. Mol. Biol. Rev.*, 64(4):655-671.
- Zimmerman S.B., Schwartz C.D., Monaghan R.L., Pleak B.A., Weissberger B., Gilfillan E.C., Mochales S., Hernandez S., Currie S.A., Tejera E. and E.O. Stapley**, 1987. Difficidin and oxydifficidin: Novel broad spectrum antibacterial antibiotics produced by *Bacillus subtilis*. *J. Antibiotics*, 40(12):1677-1681.