

# The Open Access Israeli Journal of Aquaculture – Bamidgeh

As from **January 2010** The Israeli Journal of Aquaculture - Bamidgeh (IJA) will be published exclusively as **an on-line Open Access (OA)** quarterly accessible by all AquacultureHub (<http://www.aquaculturehub.org>) members and registered individuals and institutions. Please visit our website (<http://siamb.org.il>) for free registration form, further information and instructions.

This transformation from a subscription printed version to an on-line OA journal, aims at supporting the concept that scientific peer-reviewed publications should be made available to all, including those with limited resources. The OA IJA does not enforce author or subscription fees and will endeavor to obtain alternative sources of income to support this policy for as long as possible.

## Editor-in-Chief

Dan Mires

## Editorial Board

**Sheenan Harpaz** Agricultural Research Organization  
Beit Dagan, Israel

**Zvi Yaron** Dept. of Zoology  
Tel Aviv University  
Tel Aviv, Israel

**Angelo Colorni** National Center for Mariculture, IOLR  
Eilat, Israel

**Rina Chakrabarti** Aqua Research Lab  
Dept. of Zoology  
University of Delhi

**Ingrid Lupatsch** Swansea University  
Singleton Park, Swansea, UK

**Jaap van Rijn** The Hebrew University  
Faculty of Agriculture  
Israel

**Spencer Malecha** Dept. of Human Nutrition, Food  
and Animal Sciences  
University of Hawaii

**Daniel Golani** The Hebrew University of Jerusalem  
Jerusalem, Israel

**Emilio Tibaldi** Udine University  
Udine, Italy

## Copy Editor

Ellen Rosenberg

Published under auspices of  
**The Society of Israeli Aquaculture and  
Marine Biotechnology (SIAMB),  
University of Hawaii at Manoa Library**

and  
**University of Hawaii Aquaculture  
Program** in association with  
**AquacultureHub**

<http://www.aquaculturehub.org>



UNIVERSITY  
of HAWAII  
MĀNOA  
LIBRARY



**AquacultureHub**  
educate • learn • share • engage

ISSN 0792 - 156X

© Israeli Journal of Aquaculture - BAMIGDEH.

PUBLISHER:  
Israeli Journal of Aquaculture - BAMIGDEH -  
Kibbutz Ein Hamifratz, Mobile Post 25210,  
ISRAEL

Phone: + 972 52 3965809

<http://siamb.org.il>

## Survival and Growth of Captive Reared Juvenile Seahorse (*Hippocampus kuda*) Fed Live Feeds and Fishmeal

M. Thangaraj<sup>1\*</sup> and A.P. Lipton<sup>2</sup>

<sup>1</sup> Marine Biotechnology Laboratory, Vizhinjam Research Center of Central Marine Fisheries Research Institute, Vizhinjam, Thiruvananthapuram, Kerala 695521, India

<sup>2</sup> Center for Marine Science and Technology, Manonmaniam Sundaranar University, Rajakkamangalam, Tamilnadu, 629502, India

(Received 25.2.08, Accepted 18.3.08)

Key words: seahorse juveniles, live feeds, growth, survival

### Abstract

Pond and aquarium keeping of seahorse (*Hippocampus kuda*) is challenging because of their feeding habits. Although seahorse is a purely active predator, we successfully raised laboratory-produced young on inert feed and evaluated the effects of readily-available cost-effective food items on their growth and survival. Two-month old laboratory bred juveniles were fed live *Artemia*, live *Mysids*, minced fishmeal, or a combination of *Artemia* and *Mysids* (1:1). Juveniles fed the combination diet reached the highest wet weight ( $1.93 \pm 0.31$ g), specific growth rate ( $2.21 \pm 0.07$ ), and survival ( $96 \pm 3.0\%$ ), significantly ( $p < 0.05$ ) higher than juveniles fed fishmeal ( $1.42 \pm 0.12$  g,  $2.08 \pm 0.03$ , and  $76 \pm 3.0\%$ , respectively). Although there were insignificant differences in growth and survival between juveniles fed the combination and those fed *Mysids* or *Artemia*, the combination diet was best. This easily available feed can be used in ornamental aquaria.

### Introduction

Seahorse culture has been the subject of recent attention by aquarists (Payne and Rippingale, 2000; Woods, 2000a,b) and policy planners (CITES, 2001). Concerns regarding overexploitation in the wild resulting in population declines and ever-increasing demand resulting in high market prices have led to interest in seahorse culture (Vincent, 1996; Lourie et al., 1999).

The Indian seahorse, *Hippocampus kuda*, is one of the heavily exploited seahorses along the Palk Bay coast of India (Lipton and Thangaraj, 2002). Aquaculture could reduce overexploitation by supplying the trade with bred seahorses. However, information on culturing protocols for most tropical Indo-Pacific species is limited (Do et al., 1998). Clear information on the technical feasibility of intensive

---

\* Corresponding author. Present address: Center for Advanced Study in Marine Biology, Annamalai University, Parangipettai, Cuddalore (DT), Tamilnadu 608502, India, e-mail: coralholder@yahoo.com

seahorse culture is essential for reducing overexploitation of wild stocks and to improve production.

One of the main troubles in establishing seahorse aquaculture is provision of sufficient quantities of nutritionally balanced live food. In their natural environment, seahorses are voracious predators on live amphipods, copepods, mysid shrimp, and Caridian juveniles (Lovett, 1969; Tipton and Bell, 1988). In captive conditions, aquarists and researchers have tried to feed seahorses cultivable live foods such as brine shrimp, copepods, and mysid shrimps (Lockyear et al., 1997; Wilson and Vincent, 1998; Hiloman-Garcia, 1999; Payne and Rippingale, 2000). Generally, seahorses are not fed inert foods such as frozen fish fry and *Mysids* because such foods are not readily accepted by them. However, year-round mass culture of live foods is difficult and costly. Therefore, seahorse culturists have tried alternate foods such as shrimp, fishmeal based diets, raw frozen mysids, and copepods (Chen, 1990; Forteach, 2000; Woods and Valentino, 2003).

In this study, growth and survival of *H. kuda* juveniles fed one of three types of live feed or fishmeal were compared.

#### Materials and Methods

Experiments were conducted in the wet laboratory of the Tuticorin Research Center of the Central Marine Fisheries Research Institute in Tuticorin, India, during February 2004. Flat-bottomed circular tubs (25 l) were stocked with randomly selected 60-day-old juveniles (standard length  $65.28 \pm 5.4$  mm, wet weight  $262.75 \pm 15.5$  mg) at a density of four juveniles per 10 l. Standard length and wet weight were measured as in Lourie et al. (1999). Care was taken to choose juveniles with minimum differences in standard length, wet weight, and condition factors within replicates at the start of the experiment (ANOVA,  $p > 0.05$ ).

In-coming filtered seawater was regulated at the rate of 2 l/h. Conditioned dead corals were provided as hold fasts. A photoperiod of 12 h light:12 h dark was maintained during the 30-day experiment. Water temperature was kept at  $26.0 \pm 0.85^\circ\text{C}$ , dissolved oxygen at

$5.20 \pm 0.24$  ppm, salinity at  $36 \pm 0.5$  ppt, and pH at  $8.00 \pm 0.26$ . Mean  $\pm$  SD water quality parameters did not vary between treatments or replicates (ANOVA,  $p > 0.05$ ). Tubers were inspected daily and excess food and excreta were siphoned out. Every five days, five individuals from each tub were randomly selected and standard length and wet weight were measured.

*Hippocampus kuda* juveniles were fed live *Artemia* ( $2.61 \pm 0.34$  mg) collected from open salt pans in Tuticorin, live *Mysids* ( $2.90 \pm 0.30$  mg) collected from shallow coastal seawater with a scoop net, fishmeal minced into small pieces using a sharp knife, or a combination of *Artemia*+*Mysids* (1:1). Juveniles were fed twice daily at 10:00 and 15:00.

Specific growth rate (SGR%) was calculated as the increase in body weight/day according to the formula  $(\log_{fbw} - \log_{ibw}/t) \times 100$ , where fbw = final body wt in mg, ibw = initial body wt in mg, and t = number of days. The condition factor (CF) was calculated for individual juveniles as (wet wt in g/length in mm)  $\times 100$ . Statistical analysis was performed by Statistica 5.0 software. Differences were considered significant when  $p < 0.05$ .

#### Results

**Growth and condition factor.** Juveniles fed *Artemia* and *Artemia*+*Mysids* attained mean lengths of  $101 \pm 9.45$  and  $100.50 \pm 8.90$  mm, respectively, although the differences were not statistically different (Fig. 1). The mean length of juveniles fed *Artemia* significantly differed from those fed *Mysids* while those fed fishmeal reached the significantly lowest mean length ( $90.50 \pm 9.20$  mm). Juveniles fed *Artemia*+*Mysids* attained the highest wet weight,  $1.93 \pm 0.31$  g, significantly higher than juveniles fed fishmeal,  $1.42 \pm 0.12$  g but similar to juveniles fed *Artemia* or *Mysids*. Juveniles fed fishmeal had a significantly lower SGR and condition factor than all other juveniles (Table 1).

**Survival.** The first mortality was observed on the fifth day of the experiment, 65 days post hatch (Fig. 2). At thirty days, juveniles fed fishmeal had a significantly lower survival than all other juveniles.




Fig. 1. Growth in (a) length and (b) weight of juvenile seahorse (*Hippocampus kuda*) fed one of four food items (means $\pm$ SD, n = 3).

Table 1. Specific growth rate (SGR), condition factor, and survival of juvenile *Hippocampus kuda* fed with one of four food items (means $\pm$ SD; n = 3).

Food item	SGR (%/day)	Condition factor	Survival (%)
<i>Artemia</i>	2.19 $\pm$ 0.07 <sup>a</sup>	1.81 $\pm$ 0.13 <sup>a</sup>	93.6 $\pm$ 3 <sup>a</sup>
<i>Mysids</i>	2.19 $\pm$ 0.06 <sup>a</sup>	1.98 $\pm$ 0.13 <sup>a</sup>	90.3 $\pm$ 3.3 <sup>a</sup>
Fishmeal	2.08 $\pm$ 0.03 <sup>b</sup>	1.56 $\pm$ 0.02 <sup>b</sup>	76.6 $\pm$ 3.2 <sup>b</sup>
<i>Artemia+Mysids</i>	2.21 $\pm$ 0.07 <sup>a</sup>	1.91 $\pm$ 0.05 <sup>a</sup>	96.7 $\pm$ 2.9 <sup>a</sup>

Means within a column sharing a common superscript do not significantly differ at  $p > 0.5$  for SGR and  $p > 0.05$  for condition factor and survival.




Fig. 2. Survival of juvenile seahorse (*Hippocampus kuda*) fed one of four food items (means $\pm$ SD, n = 3).

## Discussion

In seahorse aquaculture practices, the use of inert and frozen food can dramatically reduce costs of materials and labor. However, the appropriate food item varies depending on the growth stage (age). Two-month-old *H. kuda* juveniles can be accustomed to frozen and freshly dead foods (pers. observation). We observed that juveniles consume more motile food than non-motile.

In the present experiment, juveniles fed *Artemia* and *Artemia+Mysids* had better growth and survival than juveniles fed *Mysids* or fishmeal. The significantly higher growth and survival in groups fed *Artemia*, *Mysids*, and *Artemia+Mysids* probably reflect the importance of active live foods. Higher growth

was recorded in juveniles of *Hippoglossus hippoglossus* and *H. abdominalis* fed live *Artemia* (Rosenlund et al., 1997; Woods and Valentino, 2000b). In natural conditions, seahorse juveniles feed on copepods, while adults feed on small benthic crustaceans such as amphipods (Do et al., 1998). Therefore, juveniles fed *Artemia*, *Mysids*, or *Artemia+Mysids* had a sea-based food item and relatively good growth.

*Hippocampus kuda* juveniles gained 28 mm in standard length when fed *Artemia* enriched with *Acetes* for 56 days (Job et al., 2002). In the present study, juveniles fed un-enriched *Artemia* gained 36 mm in only 30 days in an ambient temperature of  $26.0 \pm 0.85^\circ\text{C}$ . The rapid growth and higher survival of *H. kuda* in our study may be due to a more appropriate temperature in Indian laboratory conditions. Thus, temperature may play a crucial role for growth and survival of young seahorse.

Frozen *Mysid* and fish-based diets would be cheaper and more convenient dietary alternatives to collection of live feeds such as *Artemia* and *Amphipods* from the wild or their maintenance in culture conditions. However, more research is required to improve diet presentation and set up suitable rearing tanks that keep non-live foods in suspension in the water column. It is also necessary to research the nutritional profile of the feeds and the nutritional requirements of *H. kuda* at different ages.

#### Acknowledgments

The authors would like to thank Prof. Dr. Mohan Joseph Modayil, Director of CMFRI, and Dr. A.C.C. Victor, Scientist-in-Charge, Dr. M. Rajamani, Dr. D.C.V. Easterson, Principal Scientists, Tuticorin R.C. of CMFR, CMFRI, Tuticorin for providing facilities. We are thankful to the Ministry of Environment and Forests, Government of India, New Delhi, for funding.

#### References

- Chen J.**, 1990. Seahorse culture. pp. 1-6. In: P. Bueno, A. Lovatelli (eds.), *Brief Introduction to Mariculture of Five Selected Species in China*. UNDP/FAO Regional Sea Forming Development and Demonstration Project, Bangkok, Thailand. National Inland Fisheries
- Institute, Kasetsart Univ., Bangkok, Bangkok, Thailand (RAS/90/002).
- CITES**, 2001. Seahorse and other members of the family Syngnathidae. Convention on International Trade in Endangered Species of Wild Fauna and Flora. Notification no. 2001/034.
- Do H.H., Truong S.K. and T.H. Ho**, 1998. Feeding behaviour and food of seahorses in Vietnam. pp. 307-319. In: B. Morton (ed.). *The Marine Biology of the South China Sea III*. Hong Kong Univ. Press, Hong Kong.
- Forteach N.**, 2000. Farmed seahorses – a boon to the aquarium trade. *Infofish Int.*, 3:48-50
- Job S.D., Do H.H., Meeuwijng J.J. and H.J. Hall**, 2002. Culturing the oceanic seahorse, *Hippocampus kuda*. *Aquaculture*, 214:333-341.
- Hiloman-Garcia G.**, 1999. AQD's marine ornamental fish project. *SEAFDEC Asian Aquac.*, 21(2):31-38.
- Lipton A.P. and M. Thangaraj**, 2002. Present status of seahorse fishing along the Palk Bay coast of Tamilnadu. *Tech. Extension Ser., Mar. Fish. Info. Serv.*, 174:5-8.
- Lockyear J., Kaiser H. and T. Hecht**, 1997. Studies on the captive breeding of the Knysna seahorse, *Hippocampus capensis*. *Aquat. Sci. Cons.*, 1:129-136.
- Lourie S.A., Vincent A.C.J. and H.J. Hall**, 1999. Seahorses: an Identification Guide to the World's Species and their Conservation. Project Seahorse, London. 214 pp.
- Lovett J.M.**, 1969. *An Introduction to the Biology of the Seahorses Hippocampus abdominalis*. B.Sc. Honors thesis, Univ. Tasmania, Australia. 102 pp.
- Payne M.F. and R.J. Rippingale**, 2000. Rearing West Australian seahorse, *Hippocampus subelongatus*, juveniles on copepod nauplii and enriched *Artemia*. *Aquaculture*, 188:353-361.
- Rosenlund G., Stoss J. and C. Talbot**, 1997. Co-feeding marine larvae with inert and live diets. *Aquaculture*, 155:183-191.
- Tipton K. and S.S. Bell**, 1988. Foraging patterns of two Syngnathidae fishes: importance of harpacticod copepods. *Mar. Ecol. Prog. Series*, 47:31-43.

- Vincent A.C.J.**, 1996. *The International Trade in Seahorses*. TRAFIC Int., Cambridge, UK. 163 pp.
- Wilson M.S and A.C.J. Vincent**, 1998. Preliminary success in closing the life cycle of exploited seahorse species, *Hippocampus* spp., in captivity. *Aquarium Sci. Conserv.*, 4: 179-196.
- Woods C.M.C.**, 2000a. Improving initial survival in cultured seahorses, *Hippocampus abdominalis* Leeson, 1827 (Teleostei: Syngnathidae). *Aquaculture*, 190:377-388.
- Woods C.M.C.**, 2000b. Preliminary observations on breeding and rearing the seahorse *Hippocampus abdominalis* (Teleostei: Syngnathidae) in captivity. *NZ J. Mar. Freshw. Res.*, 34:475-485.
- Woods C.M.C. and F. Valentino**, 2003. Frozen mysids as an alternative to live *Artemia* in culturing seahorses *Hippocampus abdominalis*. *Aquac. Res.*, 34:757-763.